



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

caught on a halibut trawl in 65 fathoms of water in Lat. 58° 5' N., Long. 149° W., a specimen of this fish which was sent to the Bureau of Fisheries by Edwin Ripley, fish dealer of Seattle, Washington. As the present specimen, which was 33 $\frac{1}{3}$  inches in total length differed somewhat from the typical description, the following notes upon it are given: Standard length 74cm. Depth, 18.5 cm. Length of caudal peduncle, 4 cm. Depth of caudal peduncle, 7.2 cm. Pectoral rounded, its longest ray 11 cm. Caudal truncate, its longest ray, 11 cm. Pectoral with small scales on base on rays, extending (on middle ray) to 6.5 cm. Dorsal, anal and caudal scaled at base. Dorsal formula, LIV. Anal, 24. Head, 13.7 cm. Eye, 2.4 cm. Snout, 3cm. Max., 4.5 cm. Interorbital, 5.3 cm. Ground color, gray, lighter beneath, punctulated with black dots. Scales with bluish margins. Dorsal and anal without punctulations. Pectoral dark at base, with dark, distal portion light gray, margin blackish. Caudal whitish at edges and towards end, after margin blackish. Pores of head, white. Forehead grayish. Yellow spot in upper axil of pectoral. B. F. tag no. 9547.

WILLIAM C. KENDALL,  
*U. S. Bureau of Fisheries.*

## COMPARATIVE NUMBERS OF LIZARDS AND SNAKES ON DESERT.

Lizards are the most abundant form of desert reptile life. They greatly outnumber the snakes both in point of individuals and species. In the summer of 1913, the writer saw only three species of snakes on the Painted Desert, Arizona. These were the prairie rattlesnake (*Crotalus confluentus*), bull snake (*Pituophis sayi*), Arizona ribbon snake(?) (*Eutaenia megalops*).

Among the lizards which are abundant may be mentioned: Bailey's Collared Lizard, Leopard Lizard, Racerunners, Swifts and Horned Lizards.

DWIGHT FRANKLIN,  
*New York, N. Y.*